

*Penstemon parryi*  
Parry's Penstemon or Beardtongue

**Why Observe?**

Parry's penstemon, also known as Parry's beardtongue, is native to Arizona. It is a hardy, drought-tolerant plant that is an important source of food for hummingbird and native bee species.

**Tips for Identification**

Will have 2-3 ft stems with clustered magenta flowers. Leaves are bluish-green, arrow-shaped, opposite, and lie mostly near the base of the plant. Flowers are tubular shaped.

Be aware there is variation from individual to individual within a species, so your plant may not look exactly like the one pictured. If you are uncertain whether or not a phenophase is occurring, report a "?" for its status until it becomes clear what you are observing after subsequent visits.



[Katja Schulz via Flickr](#), CC BY 2.0 (cropped)



[William Herron via Flickr](#), CC BY-SA 2.0

*Penstemon parryi*  
Parry's Penstemon or Beardtongue

**Image Needed**

email  
[photos@usanpn.org](mailto:photos@usanpn.org)

**Initial growth**

New growth of the plant is visible after a period of no growth (winter or drought), either from above-ground buds with green tips, or new green or white shoots breaking through the soil surface. Growth is considered "initial" on each bud or shoot until the first leaf has fully unfolded. For seedlings, "initial" growth includes the presence of the one or two small, round or elongated leaves (cotyledons) before the first true leaf has unfolded.



S Brewer. [CC BY 2.0](#)

**Leaves**

One or more live, fully unfolded leaves are visible on the plant. For seedlings, consider only true leaves and do not count the one or two small, round or elongated leaves (cotyledons) that are found on the stem almost immediately after the seedling germinates. Do not include fully dried or dead leaves.

**Flowers or flower buds**

One or more fresh open or unopened flowers or flower buds are visible on the plant. Include flower buds or inflorescences that are swelling or expanding, but do not include those that are tightly closed and not actively growing (dormant). Also do not include wilted or dried flowers.



[Katja Schulz via Flickr.](#)  
[CC BY 2.0](#)



[Kimberly Kling via Flickr.](#)  
[CC BY-SA 2.0](#)

**Open flowers**

One or more open, fresh flowers are visible on the plant. Flowers are considered "open" when the reproductive parts (male stamens or female pistils) are visible between or within unfolded or open flower parts (petals, floral tubes or sepals). Do not include wilted or dried flowers.

**Fruits**

One or more fruits are visible on the plant. For *Penstemon parryi*, the fruit is a small capsule that changes from green to tan or light brown and splits open to expose the seeds. Do not include empty capsules that have already dropped all of their seeds.



[PM Consigny via Smugmug.](#)  
[CC BY-NC-SA 2.0](#) (cropped)

**Image Needed**

email  
[photos@usanpn.org](mailto:photos@usanpn.org)

**Ripe fruits**

One or more ripe fruits are visible on the plant. For *Penstemon parryi*, a fruit is considered ripe when it has turned tan or light brown and has split open to expose the seeds. Do not include empty capsules that have already dropped all of their seeds.

**Phenophases not pictured:** Recent fruit or seed drop